

THE PREVALENCE OF CHRONIC UNTREATED HEPATITIS C VIRUS INFECTION AND ASSOCIATED RISK FACTORS IN IRISH PRISONERS; A CROSS SECTIONAL STUDY

Crowley D^{1,2}, Cullen W³, Keevans M¹, Kelly E¹, Laird E⁴, Mc Kiernan S⁶, Miggin S⁶, Mc Hugh T⁵, Murphy C^{1,5}, Murtagh R^{3,5}, O'Reilly D¹, Betts- Symonds G⁷, Tobin C⁸, Van Hout M⁹, Lambert JS^{3,5}

¹Irish Prison Service, Mountjoy Prison, Dublin, Ireland, ²Health Service Executive, Addiction Services, Dublin, Ireland, ³UCD School of Medicine, University College Dublin, Dublin, Ireland, ⁴Trinity College Dublin, Ireland, ⁵Mater Misericordiae University Hospital, Dublin, Ireland, ⁶St James Hospital, Dublin, Ireland, ⁷Irish Red Cross, Mountjoy Prison Campus, Dublin, Ireland, ⁸Psychology Department, University of Florida, USA, ⁹Public Health Institute, Liverpool John Moores University, Liverpool, UK.

Background:

An estimated 26% of prisoners globally have been exposed to hepatitis C virus (HCV) infection, increasing to 64% in those with a history of injecting drug use (IDU). There is limited data on the prevalence of hepatitis C in prisoners. This study aims to estimate the prevalence of untreated chronic HCV infection and identify associated risk factors in an Irish prison setting.

Methods:

A single site cross sectional prevalence study. Mountjoy prison – a high security male custodial prison, housing 538 male prisoners located in Dublin, Ireland. A total of 78% of prisoners consented to participate in the study, with 403 completing an enhanced HCV screening programme and 298 completing a risk questionnaire.

Results:

The prevalence estimate for untreated chronic HCV infection was 13.1%. Of those screened, 22.8 % were HCV antibody positive with 25.0 % of this group showing spontaneous clearance. Of those with chronic infection, 23.2 % had an SVR post treatment and 76.8 % untreated active HCV infection, reflecting an untreated chronic HCV infection population prevalence of 13.1%.

Of those exposed to HCV, 11% were co-infected with HIV and 6% with HBV. The prevalence of HCV infection among prisoners with a history of IDU was 79.7 %. Significant risk factors identified were IDU ($p < 0.0001$), history of prison tattoo ($p < 0.0001$), unsterile community tattoo ($p < 0.0001$), sharing needles ($p < 0.0001$) and other drug taking paraphernalia ($p < 0.0001$). The sharing of razors ($p = 0.95$) and tooth brushes ($p = 0.32$) while incarcerated was not shown to be a significant risk. On multivariate analysis, a history of non-sterile community tattoo was the only significant risk factor independent of IDU ($p = 0.005$, $\beta = 0.468$).

Conclusion:

Estimating the prevalence of untreated HCV infection in prison populations is essential to reliably inform HCV treatment strategies and to capture treatment uptake and outcomes along with HCV reinfection rates and prevention strategies.

Disclosure of Interest Statement:

HepCare Europe is funded through the European Union with partial funding received from the Health Services Executive. Dr John S Lambert has also received unrestricted grants from the Pharmaceutical Industry to support his research work.