

HEPATITIS C TREATMENT: PEER INSIGHTS TO UNDERSTANDING ENABLERS AND IMPEDIMENTS TO DAA UPTAKE AMONG PEOPLE WHO INJECT DRUGS

Chong S¹, Brown G¹

¹ Australian Research Centre in Sex, Health and Society, School of Psychology and Public Health, La Trobe University, Melbourne, Australia

Background : The listing of direct-acting antiviral (DAA) medicines for people with hepatitis C on the Australian universal health system has set the trajectory to eliminate hepatitis C. A key component is treatment for people who inject drugs (PWID) living with hepatitis C. Initial uptake of treatment among PWID was promising but uptake is now slowing. This study investigated the evolving barriers and motivators for PWID to access DAA and the implications for scaling-up testing and treatment.

Methods: PWID peer workers have unique insight into the experiences of PWID who may not participate in research/consultations. This study drew on the perspectives of peer workers and volunteers and their networks to investigate experiences of PWID regarding DAA treatment scale-up.

Ten focus groups and four semi-structured interviews were conducted across Australia in three phases with:

- I. Peer workers and volunteers from three collaborating peer-organisations to obtain insights on PWID uptake of DAA treatment
- II. individual PWID on their engagement with DAA treatment

Results: Three short reports to facilitate real-time translation of results into ongoing treatment scale-up strategies and practice were produced. Common themes included attitudes on DAAs, hepatitis C testing, access to health providers, and structural barriers to accessing treatment. We found generalized hepatitis C treatment awareness campaigns were having reduced impact, and conflicting advice and messaging, and stigmatising experiences encountered by PWID were undermining scale-up programs. Targeted strategies emphasising benefits and relevance of treatment at the individual level are needed. We also identified practices of health services demonstrating progress in engaging PWID in DAA treatment.

Conclusions: PWID peer workers and volunteers insights are vital to the development of approaches to increase access to DAA treatment. Harm reduction and hepatitis C agencies have integrated study findings to strengthen policy advocacy strategies and health promotion interventions to tailor to communities of people who inject.